

# The Relationship of Agmatine Deficiency with the Lung Cancer

Serkan KAPANCIK<sup>1</sup>, V. Kenan CELIK<sup>1</sup>, Saadettin KILICKAP<sup>2</sup>,  
Turgut KACAN<sup>3</sup>, Sercan KAPANCIK<sup>1</sup>

<sup>1</sup> Cumhuriyet University School of Medicine, Department of Biochemistry, Sivas

<sup>2</sup> Hacettepe University School of Medicine, Department of Preventive Oncology, Ankara

<sup>3</sup> Afyonkarahisar State Hospital, Department of Medical Oncology, Afyon, TURKEY

## ABSTRACT

The polyamines are essential for rapidly growing cells. Polyamines is synthesized by two reactions which begin from arginine. The aim of this study is to examine the serum levels of the enzymes located in the putrescine synthesis pathway and their relationship with lung cancer. In addition, the study also aim determine the oxidant-antioxidant status and the oxidative stress index. This study was performed on 36 lung cancer patients and 36 healthy controls. The arginase activity and the ornithine level were determined spectrophotometrically while the arginine decarboxylase, ornithine decarboxylase and agmatinase levels were determined by using Enzyme Linked Immunosorbent Assay (Elisa kit). The total antioxidant (TAS) and total oxidant status (TOS) were determined using a commercial kit auto analyser, oxidative stress index (OSI) was subsequently estimated. We observed an increase in ornithine decarboxylase, agmatinase and arginase levels in cancer patients ( $p < 0.05$ ). however, no significant difference was observed between the two groups with regards to orithine and arginase decarboxylase levels ( $p > 0.05$ ). In the patient group, the TAS values decreased ( $p < 0.05$ ), while the TOS values increased ( $p < 0.05$ ). In addition, the OSI values increased ( $p < 0.05$ ). Reduced agmatine level can cause an increase in polyamine synthesis and cell proliferation. For this reason, lung cancer prognosis may have a relationship with the agmantine levels.

**Keywords:** Lung Cancer, Polyamine, Arginase, Ornithine, Arginine decarboxylase, Ornithine decarboxylase, Agmatinase

## ÖZET

### Agmatin Eksikliğinin Akciğer Kanseriyle İlişkisi

Poliaminler, hızlı bir şekilde büyüyen hücrelerde esansiyeldir. Poliaminlerin oluşumu, arjininden başlayan iki farklı reaksiyon ile gerçekleşir. Bu çalışmadaki amaç, putresin sentez yolunda görev alan enzimlerin serum düzeylerini incelemek ve akciğer kanseri ile olan ilişkisini ortaya koymaktır. Ayrıca, oksidan-antioksidan kapasite ve oksidatif stress indeksini saptamaktır. Bu çalışmaya 36 akciğer kanseri hastası ve 36 sağlıklı kontrol dahil edildi. Arjinaz aktivitesi ve ornitin düzeyi spektrofotometrik olarak ve arjinin dekarboksilaz, ornitin dekarboksilaz, agmatinaz düzeyleri ise elisa kiti kullanılarak tayin edildi. Total antioksidan (TAS) ve total oksidan kapasite (TOS) ticari kit kullanılarak otoanalizörde saptandı. Ve oksidatif stress indeksi (OSI) hesaplandı. Çalışmada akciğer kanserli hastaların serumlarında ornitin dekarboksilaz, agmatinaz ve arjinaz düzeylerinde artış gözlemlendi ( $p < 0.05$ ). Ornitin ve arjinin dekarboksilaz düzeyleri bakımından iki grup arasında istatistiksel olarak anlamlı bir fark saptanmamıştır ( $p > 0.05$ ). Hasta grubunda TAS değerleri düştü ( $p < 0.05$ ), TOS değerleri arttı ( $p < 0.05$ ). Buna bağlı olarak OSI değerleri de arttı ( $p < 0.05$ ). Azalmış agmatin düzeyi, poliamin sentezinde ve proliferasyonda artışa neden olabilir. Bundan dolayı akciğer kanserinin prognozu ile agmatin düzeyleri birebir ilişkili olabilir.

**Anahtar Kelimeler:** Akciğer Kanseri, Poliamin, Arjinaz, Ornitin, Arjinin Dekarboksilaz, Ornitin Dekarboksilaz, Agmatinaz

## INTRODUCTION

Lung cancer is the most prevalent of cancer deaths in the world (1/5th of the deaths realised due to cancer, is from lung cancer). Due to the occurrence of approximately 1.8 million new lung cancer cases per year, it is noted to be the most prevalent cancer type.<sup>1</sup> Therefore, understanding the mechanisms of lung cancer, is important in the treatment of this disease.

The accumulation of polyamines in the tissues increase the proliferation of cells and even has a contribution in bacterial proliferation. In recent years, as a result of the aforementioned properties of polyamines, they have become a subject of research with regards to neoplastic tissues.<sup>2</sup>

The polyamines play an important role in many different biological processes such as the regulation of ion channels, enzyme activities, translation and transcription. At the same time, the polyamines constitute a response to oxidative stress and increase the mRNA levels which are necessary for protein biosynthesis by forming hypusine.<sup>3-6</sup> The regulation of the polyamine levels in the tumour cells are frequently disrupted.<sup>5,7</sup> The polyamine levels are found to be particularly higher in rapidly proliferating cells.<sup>5,7,8</sup> In studies that have been made, it was reported that the level of polyamine frequently increased in the urine and blood of breast, colon, lung, prostate and skin cancer patients. In a previous analyses carried out on cancer patients, it was observed that the major polyamines (putrescine, spermidine, spermine) and the acetylated forms of these were increased and accumulated.<sup>9-15</sup> In addition, the relationship between the inhibition of polyamine synthesis and cancer therapy are currently being investigated.<sup>10</sup> It was shown that polyamine level and the tumour size reduced with the therapy.<sup>16</sup> This indicates that cancer cells need polyamines for growth and proliferation. For this reason, it is thought that increasing polyamines provide considerable contributions to poor prognosis of cancer.<sup>17-20</sup>

In this study, we aimed to determine the levels of arginine decarboxylase, agmatinase (it gives information about the agmatine level indirectly) from the enzymes which take charge in the synthesis of putrescine that is necessary for the synthesis of the major polyamines (spermidine, spermine), argin-

ase, ornithine decarboxylase and ornithine in the serums of lung cancer patients and to put forth its relationship with lung cancer. In addition, to examine the relationship between these parameters and oxidative stress factors.

## MATERIALS AND METHODS

### Patient and Control Group

The 36 patients, who were diagnosed with lung cancer and applied to Cumhuriyet University Medicine Faculty Oncology Polyclinic were included in the study. No restriction was made with respect to sex, age or the histopathologic type of cancer as well as its degree.

Our control group consisted of 36 healthy individuals who had no systemic diseases (diabetes, hypertension, lung cancer disease).

### Blood Samples Collection

Blood samples were taken from the controls and lung cancer cases who were not on therapy. The serums were obtained by centrifugation at 1610 x g for 10 minutes from the blood sample preserved at -80°C for the study.

Ornithine, arginine decarboxylase, ornithine decarboxylase, agmatinase levels and arginase enzyme activity have been measured in serum of lung cancer patients.

Arginase activity were measured spectrophotometrically according to the method defined by Geyer and Dabich and was reported as  $\mu\text{mol urea/mL/hour}$ .<sup>21</sup> Ornithine level were calculated spectrophotometrically at 515 nm using the method defined by Chinard and was reported as  $\mu\text{mol/mL}$ .<sup>22</sup>

Arginine decarboxylase, ornithine decarboxylase and agmatinase levels were determined by ELISA kit according to the manufacturer's protocol (Cusabio Biotech, Wuhan, China).

The total antioxidant and the total oxidant status were determined by using commercial kit (Rel Assay Diagnostic)

### Statistical Analysis

In the evaluation of data, when parametric test assumptions were fulfilled the significance test of the

**Table 1.** The age ranges of lung cancer patients and controls

Age (in Years)	Patient (n)	Control (n)	P
0-50	4	7	
51-60	11	19	
61-70	15	7	
>71	6	3	
* Median age	61,72±9,04	57,44±15,51	P= 0.157
* Data expressed as mean ± standard deviation			

difference between the two average was applied, the parametric test assumptions were not fulfilled the Mann Whitney U test was applied. The data was given in the tables as arithmetical average± Standard deviation and the level of significance was taken as less than 0,05.

## RESULTS

When the ages of the individuals in the control group (mean age  $57.44 \pm 15.51$ ) were compared with the ages of the individuals in the case group (mean age  $61.72 \pm 9.04$ ), there was no significant difference observed ( $p= 0.157$ ) (Table 1). Also, there was no significant difference observed between gender in the case and control group.

The tumor characteristics of lung cancer patients were given on Table 2.

In our study, when individuals in the case and control groups were compared in respect to their levels of agmatinase, arginase and ornithine decarboxylase, the difference between the two groups was

**Table 2.** The tumor characteristics of lung cancer patients

Tumor Histology and Stage	Patient (n)
Adenocarcinoma	
Localised	0
Locally Advanced	3
Metastatic	8
Squamous cell carcinoma	
Localised	0
Locally Advanced	11
Metastatic	7
Small cell lung cancer	
Localised	0
Locally Advanced	1
Metastatic	4
Non-small cell lung cancer	
Localised	0
Locally Advanced	1
Metastatic	1

statistically significant ( $p= 0.001$ ) The increase in the enzyme levels were compared in accordance to the control group, respectively agmatinase was 300%, arginase was approximately 200% and ornithine decarboxylase was 100%. Statistically, there was no significant difference between the two groups with regard to their ornithine and arginine decarboxylase levels(respectively,  $p= 0.860$  and  $p= 0.424$ ) (Table 3).

Also in our study, when individuals in the case and control groups were compared with regard to total antioxidant status, total oxidant status and oxidative stress index, the difference between the two groups was statistically significant (respectively,  $p= 0.002$ ,  $p= 0.001$  and  $p = 0.001$ ) When these pa-

**Table 3.** The serum Ornithine, Arginine Decarboxylase, Ornithine Decarboxylase, Agmatinase levels and Arginase activity of lung cancer patients and controls

	Patient ( n=36 )	Control ( n=36 )	p
**Arginine Decarboxylase (pg/ml)	206.36±109.66	191.01±126.67	0.424
**Ornithine Decarboxylase (pg/ml)	225.47±122.40	130.29±70.42	0.001
**Agmatinase (pg/ml)	967.72±769.89	233.70±109.88	0.001
**Arginase (µmol urea/ml/hour)	23.88±19.58	7.38±3.91	0.001
*Ornithine (µmol/ml)	0.08±0.02	0.08±0.02	0.860
Data expressed as mean ± standard deviation			
* The significance test of the difference between the two averages.			
**Mann Whitney U test was applied.			

**Table 4.** The serum Total Antioxidant Status, Total Oxidant Status, Total Oxidative Index of lung cancer patients and controls

	Patient ( n=36 )	Control ( n=36 )	p
*Total Antioxidant Status (mmol/Trolox Equiv./L)	2.20±0.53	2.57±0.45	0.002
**Toplam Oxidant Status (µmol H <sub>2</sub> O <sub>2</sub> Equiv./L)	50.93±29.72	28.61±14.61	0.001
**Oxidative Stress Index	22.72±12.32	11.76±9.04	0.001
Data expressed as mean ± standard deviation			
* The significance test of the difference between the two averages.			
** Mann Whitney U test was applied.			

rameters were compared according to the control group, respectively, while the total antioxidant status was diminished by 10%, the total oxidant status increased by 100%. This case caused an increase of 100% in the oxidative stress index (Table 4).

## DISCUSSION

It is known that there are rapid growth in the cells and tissues which have polyamine accumulation. For this reason, it plays an important role in the formation of benign and malign disease. The polyamines are necessary for protein synthesis and response to oxidative stress which exist in many diseases. The polyamine regulation is disrupted due to the enzymatic changes in cancer patients and this lead to an increase in polyamine levels.<sup>4,10,23</sup>

In this study, when the levels of arginase and ornithine decarboxylase from the enzymes which take place in the first reaction mechanism of the putrescine formation pathway were compared according to the control group, the result revealed an increase of 200% and 100% respectively. This increase is significant statistically ( $p < 0.05$ ). In various studies that were made, it was reported that arginase activity increased different cancer types (lung, colorectal, prostate, pancreas, skin and stomach cancer).<sup>24-29</sup> The findings in these studies support our findings. The increased arginase activity must have caused an increase in ornithine levels according to the reaction. However, in our findings, it was determined that ornithine levels did not change when compared with the control group ( $p > 0.05$ ). The reason for this was the increase in ornithine decar-

boxylase level which took place in the continuation of the reaction. Thus, it demonstrated that ornithine which formed by the increased arginase activity changed into putrescine. for this reason, there was no significant difference observed when the ornithine level was compared between two group.

In a study, it was reported that when ornithine decarboxylase expression on cancerous lung and the normal lung tissue was examined, the expression of the enzyme on the cancerous lung tissue increased.<sup>30</sup> Also, it was shown that the ornithine decarboxylase expression in the other cancerous tissues such as prostate, breast, oesophagus cancer increase.<sup>31-33</sup> Besides, the increase of the mucosal ornithine decarboxylase enzyme activity in colorectal cancer and stomach cancer support our findings.<sup>34,35</sup>

When the enzyme levels in the second pathway taking place in the putrescine formation were compared to the control group, it was found that arginine decarboxylase levels did not change ( $p > 0.05$ ), but the agmatinase level increased with a ratio of 300% ( $p < 0.05$ ). In this case, the putrescine formation increase. On the other hand, the enzyme level's raising to 4 times should be diminish the amount of agmatine which is a natural substrate of the enzyme.

In the researches that were made, it was found that agmatine inhibited cell growth and proliferation.<sup>36,37</sup> It was also fixed that agmatine decreased polyamine synthesis by increasing the expression of antizyme that mediates the degradation of ornithine decarboxylase.<sup>36</sup> Another effect of the agmatine on the polyamine levels was also the diminishing of putrescine, spermidine and spermine

in the cell by increasing the spermidine-spermine acetyl transferase (SSAT) activity.<sup>38</sup>

The increase in polyamine concentration causes the increase of growth stimulation by means of increase of the transcription factors such as c-myc, c-fos and c-jun expression, the decrease in polyamine level arrest the improvement of the cell-cycle.<sup>39-41</sup> The decreasing polyamine levels stabilize the transcripts (p53, TGF- $\beta$  ve junD) of the genes which inhibited growth.<sup>42-44</sup> The absence of spermidine causes inhibition of the hipusination. This case causes inhibition of the function of factor eukaryotic translation initiation factor 5A (eIF5A) which is bounded to the hipusination and a decrease of the cell growth.<sup>45,46</sup> The polyamine levels in different cancer patients found is high compared to the control group.<sup>47-48</sup> These data strongly support our finding.

Oxidative stress can be defined as an increase in the oxidant molecules or a decrease in the antioxidant capacity. The increasing oxidative stress index causes the decrease in the defense capability of the organism against the oxidative damage.<sup>49</sup> In our study, when the oxidative stress index was compared with the control group, a 100% increase in the patient group was determined. The polyamines owned a capability of forming a response against the oxidative damage.<sup>50</sup> According to our findings, the increasing polyamine levels did not prevent the oxidative damage.

In conclusion, the findings obtained from this pilot study suggested that the increase in the enzyme levels which take part in the putrescine formation pathway causes more putrescine by influencing the direction of the synthesis. The increasing putrescine concentration naturally cause an increase in the synthesis of spermidine and spermine from the other major polyamines. While the increase in the polyamine levels inevitably causes the uncontrolled proliferation and increasing prevalence of the cancer cells, it fails in the prevention of the oxidative damage.

The agmatine can prevent the excessive putrescine formation in the synthesis reaction mechanism. It accomplishes this through two way (its effect on the two critical enzymes). In the first, agmatine inhibit the expression of the ornithine decarboxylase enzyme in the first synthesis pathway by means of

antizyme molecule. In the second, it decreases the putrescine, spermidine and spermine levels by increasing the SSAT activity.

## Conclusions

The increase of polyamine synthesis which play a role in the progression of lung cancer stem from agmatine inadequacy resulting from high agmatinase levels together with the differentiated enzyme levels in the polyamine biosynthesis pathway. The reason for this is non-prevention of the increase in the amount of putrescine which is a precursor in the synthesis of other major polyamines because of remain incapable the decreased agmatine's regulating role on the putrescine formation mechanism. There is a need for studying the relationship between agmatine and cancer prognosis in other types of cancer, as well. Future studies on this subject may provide greater contribution to cancer treatment.

## ACKNOWLEDGEMENTS

*This study was supported by Cumhuriyet University, The Scientific Research Projects Commission. (Project No. T-540)*

## REFERENCES

1. [http://globocan.iarc.fr/Pages/fact\\_sheets\\_cancer.aspx](http://globocan.iarc.fr/Pages/fact_sheets_cancer.aspx) (Access date: 01/22/2014).
2. Bachrach U. Polyamines and cancer: minireview article. *Amino Acids* 26: 307-309, 2004.
3. Park MH, Joe YA, Kang KR, et al. The polyamine-derived amino acid hypusine: its posttranslational formation in eIF-5A and its role in cell proliferation. *Amino Acids* 10: 109-121, 1996.
4. Pignatti C, Tantini B, Stefanelli C, Flamigni F. Signal transduction pathways linking polyamines to apoptosis. *Amino Acids* 27: 359-365, 2004.
5. Wallace HM, Fraser AV, Hughes A. A perspective of polyamine metabolism. *Biochem J* 376: 1-14, 2003.
6. Wang JY, Casero RA. Polyamine cell signaling: physiology, pharmacology, and cancer research. 1st edition. New Jersey, Humana Press, 2006: 349-383.
7. Nowotarski SL, Woster PM, Casero RA Jr. Polyamines and cancer: implications for chemotherapy and chemoprevention. *Expert Rev Mol Med* 15: 1-21, 2013.
8. Porter CW, Bergeron RJ. Spermidine requirement for cell proliferation in eukaryotic cells: structural specificity and quantitation. *Science* 219: 1083-1085, 1983.



9. Russell DH, Gullino PM, Marton LJ, LeGendre SM. Polyamine depletion of the MTW9 mammary tumor and subsequent elevation of spermidine in the sera of tumor-bearing rats as a biochemical marker of tumor regression. *Cancer Research* 34: 2378-2381, 1974.
10. Gerner EW, Meyskens FL. Polyamines and cancer: old molecules, new understanding. *Nature Reviews Cancer* 4: 781-792, 2004.
11. Thomas T, Thomas TJ. Polyamine metabolism and cancer. *Journal of Cellular and Molecular Medicine* 7: 113-126, 2003.
12. Watanabe S, Sato S, Nagase S, Ohkuma S. Chemotherapeutic choice of ranimustine or nimustine on the basis of regional polyamine levels in rat brain. *Methods and Findings in Experimental and Clinical Pharmacology* 30: 115-120, 2008.
13. Roy UKB, Rial NS, Kachel KL, Gerner EW. Activated K-RAS increases polyamine uptake in human colon cancer cells through modulation of caveolar endocytosis. *Molecular Carcinogenesis* 47: 538-553, 2008.
14. Byun JA, Lee SH, Jung BH, et al. Analysis of polyamines as carbamoyl derivatives in urine and serum by liquid chromatography-tandem mass spectrometry. *Biomedical Chromatography* 22: 73-80, 2008.
15. Giskeødegård GF, Bertilsson H, Selnæs KM, et al. Spermine and citrate as metabolic biomarkers for assessing prostate cancer aggressiveness. *PLoS One* 8: 1-9, 2013.
16. Verma AK. Inhibition of tumor promotion by DL- $\alpha$ -difluoromethylornithine, a specific irreversible inhibitor of ornithine decarboxylase. *Basic Life Sci* 52: 195-204, 1990.
17. Löser C, Fölsch UR, Paprotny C, Creutzfeldt W. Polyamines in colorectal cancer. Evaluation of polyamine concentrations in the colon tissue, serum, and urine of 50 patients with colorectal cancer. *Cancer* 65: 958-966, 1990.
18. Chatel M, Darcel F, Quemener V, et al. Red blood cell polyamines as biochemical markers of supratentorial malignant gliomas. *Anticancer Res* 7: 33-38, 1987.
19. Kubota S, Okada M, Yoshimoto M, et al. Urinary polyamines as a tumor marker. *Cancer Detect Prev* 8: 189-192, 1985.
20. Uehara N, Shirakawa S, Uchino H, Saeki Y. Elevated contents of spermidine and spermine in the erythrocytes of cancer patients. *Cancer* 45: 108-111, 1980.
21. Geyer JW, Dabich D. Rapid method for determination of arginase activity in tissue homogenates. *Anal Biochem* 39: 412-417, 1971.
22. Chinard FP. Photometric Estimation of Proline and Ornithine. *J Biol Chem* 199: 91-95, 1952.
23. LaMuraglia GM, Lacaine F, Malt RA. High ornithine decarboxylase activity and polyamine levels in human colorectal neoplasia. *Ann Surg* 204: 89-93, 1986.
24. Gökmen SS, Yörük Y, Çakir E, et al. Arginase and ornithine, as markers in human nonsmall cell lung carcinoma. *Canc Biochem Biophys* 17: 125-131, 1999.
25. Leu SY, Wang SR. Clinical significance of arginase in colorectal cancer. *Cancer* 70: 733-736, 1992.
26. Harris BE, Pretlow TP, Bradley ELJ, et al. Arginase activity in prostatic tissue of patients with benign prostatic hyperplasia and prostatic carcinoma. *Cancer Res* 43: 3008-3012, 1983.
27. Löser C, Fölsch UR, Paprotny C, Creutzfeldt W. Polyamine concentrations in pancreatic tissue, serum, and urine of patients with pancreatic cancer. *Pancreas* 5: 119-127, 1990.
28. Gökmen SS, Aygit AC, Ayhan MS, et al. Significance of arginase and ornithine in malignant tumors of the human skin. *J Lab Clin Med* 137: 340-344, 2001.
29. Wu CW, Chi CW, Ho CK, et al. Effect of arginase on splenic killer cell activity in patients with gastric cancer. *Dig Dis Sci* 39: 1107-1112, 1994.
30. Tian Y, Huang Q, Li L, et al. Gene expression of ornithine decarboxylase in lung cancers and its clinical significance. *Acta Biochimica et Biophysica Sinica* 38: 639-645, 2006.
31. Mohan RR, Challa A, Gupta S, et al. Overexpression of ornithine decarboxylase in prostate cancer and prostatic fluid in humans. *Clinical Cancer Research* 5: 143-147, 1999.
32. Deng W, Jiang X, Mei Y, et al. Role of ornithine decarboxylase in breast cancer. *Acta Biochimica et Biophysica Sinica* 40: 235-243, 2008.
33. Yoshida M, Hayashi H, Taira M, Isono K. Elevated expression of the ornithine decarboxylase gene in human esophageal cancer. *Cancer Research* 52: 6671-6675, 1992.
34. Hixson LJ, Garewal HS, McGee DL, et al. Ornithine decarboxylase and polyamines in colorectal neoplasia and mucosa. *Cancer Epidemiology Biomarkers and Prevention* 2: 369-374, 1993.
35. Okuzumi J, Yamane T, Kitao Y, et al. Increased mucosal ornithine decarboxylase activity in human gastric cancer. *Cancer Research* 51: 1448-1451, 1991.
36. Satriano J, Matsufuji S, Murakami Y, et al. Agmatine suppresses proliferation by frameshift induction of antizyme and attenuation of cellular polyamine levels. *J Biol Chem* 273: 15313-15316, 1998.
37. Regunathan S, Youngson C, Raasch W, et al. Imidazoline receptors and agmatine in blood vessels: a novel system inhibiting vascular smooth muscle proliferation. *J Pharmacol Exp Ther* 276: 1272-1282, 1996.
38. Vargiu C, Cabella C, Belliardo S, et al. Agmatine modulates polyamine content in hepatocytes by inducing spermidine/spermine acetyltransferase. *Eur J Biochem* 259: 933-938, 1999.
39. Wang JY, McCormack SA, Viar MJ, et al. Decreased expression of protooncogenes c-fos, c-myc, and c-jun following polyamine depletion in IEC-6 cells. *Am J Physiol Gastrointest Liver Physiol* 265: 331-338, 1993.
40. Li L, Rao JN, Bass BL, Wang JY. NF-kappaB activation and susceptibility to apoptosis after polyamine depletion in intestinal epithelial cells. *Am J Physiol Gastrointest Liver Physiol* 280: 992-1004, 2001.
41. Stephenson AH, Christian JF, Seidel ER. Polyamines regulate eukaryotic initiation factor 4E binding protein 1 gene transcription. *Biochem Biophys Res Commun* 323: 204-212, 2004.

42. Patel AR, Wang JY. Polyamines modulate transcription but not posttranscription of c myc and cjun in IEC-6 cells. *Am J Physiol* 273: 1020-1029, 1997.
43. Li L, Liu L, Rao JN, et al. JunD stabilization results in inhibition of normal intestinal epithelial cell growth through p21 after polyamine depletion. *Gastroenterology* 123: 764-779, 2002.
44. Li L, Li J, Rao JN, et al. Inhibition of polyamine synthesis induces p53 gene expression but not apoptosis. *Am J Physiol* 276: 946-954, 1999.
45. Schnier J, Schwelberger HG, Smit-McBride Z, et al. Translation initiation factor 5A and its hypusine modification are essential for cell viability in the yeast *Saccharomyces cerevisiae*. *Mol Cell Biol* 11: 3105-3114, 1991.
46. Zuk D, Jacobson A. A single amino acid substitution in yeast eIF-5A results in mRNA stabilization. *EMBO J* 17: 2914-2925, 1998.
47. Horn Y, Beal SL, Walach N, et al. Relationship of urinary polyamines to tumor activity and tumor volume in patients. *Cancer Research* 44: 4675-4678, 1984.
48. Kalac P. Health effects and occurrence of dietary polyamines: A review for the period 2005–mid 2013. *Food Chemistry* 161: 27-39, 2014.
49. Cete S, Arslan F, Yasar A. Aloe vera ve nerium oleanderin bazı mikroorganizmalara karsi antimikrobiyal aktivitenin arastirilmesi ve bu bitkilerin siklosporinli karaciger dokusundaki ksantin oksidaz enzim aktivitesine etkilerinin incelenmesi. *G Ü Fen Bilimleri Dergisi* 18: 375-380, 2005.
50. Tkachenko A, Nesterova L, Pshenichnov M. The role of the natural polyamine putrescine in defense against oxidative stress in *Escherichia coli*. *Arch Microbiol* 176: 155-157, 2001.

#### Correspondence

Serkan KAPANCIK  
Cumhuriyet Üniversitesi Tıp Fakültesi  
Biyokimya Anabilim Dalı  
58140, SIVAS / TURKEY

Tel: (+90-346) 219 10 10 / 1075  
email: serkankapancik@gmail.com